

High Frequency of Fetal Loss in Fetuses With Normal Karyotype and Nuchal Translucency ≥ 3 Among the Iranian Pregnant Women

Shokoh Abotorabi; M.D.¹, Niloufar Moeini; M.Sc.², Sahar Moghbelinejad; Ph.D.^{3,4}

¹ Department of Obstetrics and Gynecology, School of Medicine, Qazvin University of Medical Science, Qazvin, Iran

² Department of Biology, School of Basic Science, Qom Islamic Azad University, Qom, Iran

³ Cellular and Molecular Research Centre, Qazvin University of Medical Sciences, Qazvin, Iran

³ Khatam Pathobiology and Genetic Lab, Qazvin, Iran

Received November 2019; Revised and accepted June 2020

Abstract

Objective: The purpose of this study was comparison of association of three main first trimester screening factors with pregnancy outcomes among Iranian pregnant women.

Materials and methods: This prospective study was done during 2017-2019 years in Qazvin, Iran. To do so, a total of 1500 pregnant women in first trimester were enrolled. At the first step, Nuchal translucency (NT) was measured in 11-13 \pm 5 week, then the serum pregnancy-associated plasma protein A (PAPP-A) and free- β -human chorionic gonadotropin (free- β -HCG) were measured in 12-14 weeks of gestation. Pregnant women were followed up until the end of pregnancy for the complications of pregnancy such as intra-uterine growth retardation (IUGR), intrauterine death (IUFD), different types of fetal loss and preterm labor.

Results: The results showed that low levels of serum biomarkers had more association with pregnancy complications in comparison to high levels of them. Significant association of IUGR ($P = 0.001$), IUFD ($P = 0.032$) and pre-term labor ($P = 0.002$) was shown in women with low serum levels of PAPP-A in comparison to low serum levels of free- β -hCG. Significant high frequency of different types of fetal loss (IUFD, Abortion, Elective termination) was shown in fetuses with $N \geq 3$ in comparison to low levels of serum biomarkers ($P = 0.001$).

Conclusion: This study highlighted the importance of accurately interpreting the results of the first trimester of pregnancy screening which should be considered by primatologists for subsequent pregnancy care.

Keywords: Free-Beta Human Gonadotropin; Pregnancy Associated Protein-A; Nuchal Translucency; Pregnancy

Introduction

Pregnancy-associated plasma protein A (PAPP-A), Free Human chorionic gonadotropin (Free- β -HCG),

and Nuchal translucency (NT) measurements in the first trimester of pregnancy are combined screening for evaluation of fetus chromosomal abnormality. Measurement of these parameters is based on the recommendations of the Fetal Medicine Foundation (FMF) (1, 2). PAPP-A is a protein associated with pregnancy which is produced by both the embryo and

Correspondence:

Dr. Sahar Moghbelinejad

Email: smoghbelinejad@qums.ac.ir

the placenta during pregnancy (1). Its main function is proteolysis of insulin-like growth factor binding protein 4 and 5 (IGFBP 4 and 5) (3). It has major role in the local proliferation. Protease activity of PAPP-A increases the availability of IGF and causes the transfusion of glucose and amino acids in the placenta (4). The amount of PAPP-A increases from the first detection in the first trimester until the term. Low level of PAPP-A (≤ 0.4 MoM) in first trimester is associated with pregnancy complications such as: intrauterine growth restriction (IUGR) of fetus, IUFD (Intra uterine fetal death), preeclampsia and preterm birth (4-6).

HCG is a member of the glycoprotein hormone (GPH) family; HCG is heterodimer protein consisting of α - and β - subunits; α -subunit is common to all GPHs, but biological activity of HCG hormone is in β subunit and distinguishes hCG from other glycoprotein hormones. HCG functions during the pregnancy are mitotic growth, differentiation of the endometrium, localized suppression of the maternal immune system, modulation of uterine morphology, gene expression and coordination of intricate signal transduction between the endometrium. B-hCG lacks hCG activity, but several lines of study indicate that it exerts growth promoting activity. It has been speculated that b-hCG interferes with the growth-inhibiting effect of transforming growth factor-b, platelet-derived growth factor-B and nerve growth factor (7, 8). The peak of maternal serum hCG is at 8–10 weeks and then reduces to reach a plateau at 18–20 weeks. The same as PAPP-A, decreasing of free β -HCG (≤ 0.5 MoM) is associated with some pregnancy complication such as preeclampsia and preterm birth and etc. (9).

Around 11th to 14th weeks of pregnancy Nuchal translucency (NT) is measured by sonography in fetus (10). The size of NT shows the amount of fluid behind the neck of the fetus. There is a significant correlation between the increased NT and the chromosomal abnormalities, specialty trisomy 21. Therefore, with other factors such as mother age, presence of nasal bone, and blood flow across the tricuspid valve or in the ductus venous detection, the rate of Down syndrome improves to 85 % in first trimester of pregnancy. The amount of fluid behind neck can be increased in fetus with normal karyotype. In this condition fetus is at a significant risk of still birth, heart defect, and delay in neurodevelopment (10-12).

According to the above contents, the aim of this study was to analyze the association of different rates of NT, serum PAPP-A, Free- β -HCG with pregnancy outcome among the Iranian pregnant women with

normal karyotype fetus.

Materials and methods

This study was performed between 2017- 2019 years and was approved by the Ethics Committee of Qazvin University of Medical Sciences (IR.QUMS.REC.1399.113). Each of the recruited patients gave their informed written consent. In this timespan, 1500 pregnant women were referred to our laboratory by perinatologist. The first trimester screening took place at $11-13 \pm 5$ weeks of pregnancy and estimation of pregnancy week was according to the results of the first trimester ultrasound. The candidate women were also informed about the limitations of the screening method.

Before measuring the level of PAPP-A and Free- β -HCG in the serum of pregnant women, measurement of NT was done in a sonography center; then based on the recommended criteria by the FMF, risks were calculated according to the FMF program (13, 14). For evaluation of serum parameters, a clotted blood sample was obtained and serum PAPP-A and free β -hCG were measured using Cobas E 411 analyzer (made by Roche company, Germany, 2011). Serum parameters were converted into multiples of the median (MoM), then adjusted for some parameters such as the size of NT, maternal age, weight, history of Down syndrome, etc. Risk assessment of first-trimester was done according to the Feto-maternal module of the Astraia software (version 1.18.088). A calculated risk of equal to or higher than 1: 250 was defined as 'high-risk', between 250 and 1500 as intermediate risk and lower than 250 as low risk. A perinatologist recommended amniocentesis or chorionic villus sampling for evaluation of chromosomal abnormality in high risk situations. A receipt of the information was obtained from all participants. Pregnant mothers with different ranges of NT, PAPP-A, free β -hCG and normal karyotype were followed up for the outcome of pregnancy by contacting their obstetricians and study of the documents.

In fetuses with high NT ($NT \geq 3$) and normal karyotype, the size of NTs were classified into 4 levels (namely 3-4, 4-5, 5-6 and ≥ 6) and the frequency of some pregnancy complication parameters such as different types of fetal loss (Abortion, IUFD, and elective termination) and heart abnormality was evaluated in these groups. Also, the pregnancy complication factors compared in two groups of pregnant women with $0.4 \leq PAPP-A$ and $0.5 \leq \beta$ -hCG and those with $PAPP-A \leq 0.4$ and β -hCG ≤ 0.5 .

Statistical analysis: The results were analyzed by

the GraphPad analytical software (GraphPad PRISM V 5.04). The association between different ranges of the first trimester screening markers and pregnancy complications was assessed by computing the odds ratio (OR) and 95% confidence intervals (95% CI) from logistic regression analyses. We also used the Hardy–Weinberg equilibrium test for evaluation of different pregnancy outcomes frequency. All P-values were two-tailed and with $P < 0.05$ considered statistically significant

Results

In this research, 1500 pregnant women were evaluated in the first trimester (1470 singleton and 30 twin pregnancies). The mean of maternal age was 31 ± 2.82 years and 21.4% of them were 35 years of age or older. Demographic characteristics of the studied pregnant women are summarized in tables 1 and 2.

Table 1: Demographic characteristics of studied population

Characters (n = 1500)	n (%)
Mother age	
17 ≤ age ≤ 20	30 (2)
20 ≤ age ≤ 30	720 (48)
30 ≤ age ≤ 40	600 (40)
≥ 40	150 (10)
Mode of conception	
Spontaneous	1350 (90)
Assisted	75 (5)
Not reported	75 (5)
Previous affected pregnancy	
Yes	120 (8)
No	1380 (92)
Gestational age at screening time	
11	300 (20)
12	870 (58)
13	33 (22)

Totally, 24% of the studied fetuses had an estimated risk equal to or more than 1:250.

Table 2: Frequency of different chromosomal abnormalities in aneuploid fetuses

Types of chromosomal abnormality	n (%)
Numerical	
Trisomy 21	30 (2)
Trisomy 18	5 (0.3)
Trisomy 13	2 (0.13)
Monosomy X	0 (0)
Triploidy	2 (0.13)
Structural	
Tr (2;6) (p13;p23)	1 (0.06)
In 8 (p12;q21)	1 (0.06)
Del X (p1.1)	1 (0.06)

Tr: Translocation, In: Inversion, Del: Deletion

Among the age, ultrasound, NT, and biochemical factors, $NT \geq 3$ had the highest detection rate of Down syndrome (66.6%) as can be seen in Table 3. In our study 42 (2.8%) of fetuses had chromosomal aberrations: trisomy 21 (n = 30), trisomy 18 (n = 5), trisomy 13 (n = 2), triploidy (n = 2), and structural chromosomal abnormality (n = 3) (Table 3).

In another part of this research, we classified PAPP-A and free β -hCG cutoff points to low, and high ranges; then, different pregnancy complications such as IUGR, different types of fetal loss (Abortion, IUFD, elective termination), and preterm labor were compared among these groups. Our evaluation results showed that $PAPP-A \leq 0.4$ and free β -hCG ≤ 0.5 had adverse effect on pregnancy outcomes in comparison to $PAPP-A \geq 2$ and free β -hCG ≥ 2 . About $PAPP-A \leq 0.4$, frequency of IUGR was 18.5% but this frequency in high range was 1.8% ($P = 0.001$). Moreover, preterm labor frequency in samples with low level PAPP-A was significantly higher than (14.8%) samples with high level PAPP-A (6.5%) ($P = 0.001$).

Regarding the different types of fetal loss (Abortion, IUFD and elective termination) frequency we did not observe significant difference between samples with low and high ranges of PAP-A (Table 4).

Table 3: Detection rate of age, ultrasound, and serum biochemical factors among the Iranian pregnant women

Characters (n = 1500)	n(%)	Risk ≥ 1:250 [n(%)]	Detection rate [n(%)]		
			Trisomy 21	Trisomy 18	Trisomy 13
Maternal age					
17 ≤ age ≤ 20	30 (2)	5	2/30: %6.6	0 (0)	0 (0)
20 ≤ age ≤ 30	720 (48)	158	8/30: %26.6	2/5: %40	1/2: %50
30 ≤ age ≤ 40	600 (40)	167	18/30: %60	3/5: %60	1/2: %50
≥ 40	150 (10)	42	2/30: %6.6	0 (0)	0 (0)
$NT \geq 3$	105 (7)	40	2/30: %6.6	3/5: %60	1/2: %50
Absent of nasal bone	15 (1)	9	13/30: %42.4	2/5: %40	0 (0)
β -hCG ≥ 2 MoM	100 (6)	21	10/30: %33.33	0 (0)	0 (0)
$PAPP-A \leq 0.5$ MoM	102 (6.8)	28	6/30: %17.8	3/5: %60	2/2: %100
Tricuspid regurgitation	9 (0.6)	5	12/30: %36.3	4/5: 80	0 (0)

MOM: Multiple of median, β -hCG: Free beta human chronic gonadotropin, PAPP-A: Pregnancy associated protein

Table 4: Comparison of frequency of various pregnancy complications between samples with the cutoff points of PAPP-A and B-HCG

	Papp-A ≤ 0.4	Papp-A > 2	OR (95% CI)	P- value	Fβ-hCG ≤ 0.5	Fβ-hCG > 2	OR (95% CI)	P- value
IUGR	277 (18.5)	27 (1.8)	2.13 (0.32-7.13)	< 0.001	73 (4.9)	47 (3.1)	1.02 (0.2-1.2)	0.07
Normal growth	1222 (81.5)	1473 (98.2)			1427 (95.1)	1453 (96.9)		
Pre-term Labor	222 (14.8)	98 (6.5)	1.76 (0.11-6.1)	< 0.001	61 (4.06)	39 (2.65)	0.78 (0.22-0.65)	0.23
Term-labor	1278 (85.2)	1402 (93.5)			1483 (98.86)	1461 (97.4)		
Different types of fetal loss								
IUFD	132 (8.8)	22. (1.5)	1.9 (0.44-2.1)	0.03	79 (5.3)	13 (2.2)	0.92 (0.2-0.87)	0.12
Normal pregnancy	1368(91.2)	1477 (98.5)			14205 (94.7)	1467 (97.8)		
Abortion	18(1.2)	5205(3.5)	0.84 (0.11-0.87)	0.089	127 (8.47)	28 (1.9)	2.92 (0.85-6.1)	0.003
Not abortion	1482(98.8)	1477 (96.5)			1372 (91.53)	1483 (98.9)		
Elective termination	34 (2.3)	7 (0.5)	0.3 (0.1-0.75)	0.12	46 (3.1)	16 (1.12)	0.12 (0.01-1.1)	0.34
Not elective termination	1465 (97.7)	1492 (99.5)			1453 (96.9)	1483 (98.9)		

P-value shows comparison results of between high and low levels of serum markers. (Logistic regression analyses test). Fβ-hCG: Free beta human chronic gonadotropin, PAPP-A: Pregnancy associated protein A, IUFD: Intra uterine fetal death, IUGR: Intra uterine growth retardation.

Comparison of association of low and high ranges of free β-hCG with different pregnancy outcomes showed no significant difference between low and high levels of free β-hCG; we only observed high frequency of abortion in women with free β-hCG ≤ 0.5 in comparison to high range of this factor (P = 0.003). The comparison of pregnancy outcomes of patients with PAPP-A ≤ 0.4 and free β-hCG ≤ 0.5 showed a high frequency of IUGR (P < 0.001), IUFD (P = 0.032), and Pre-term labor (P = 0.002) in samples with PAPP-A ≤ 0.4 when compared to pregnancies with free β-hCG ≤ 0.5; nevertheless, we observed high frequency of abortion in samples with free β-hCG ≤ 0.5 in comparison to PAPP-A ≤ 0.4 (P = 0.001) (Table 4). In the studied samples, 105 of the fetuses had NT ≥ 3 and 38% of them had Down syndrome risk ≥ 250. We observed high frequency of abortion (P < 0.001), IUFD (P < 0.001) and elective termination of pregnancy (P < 0.001) in the samples with high NT in comparison to normal NT (Table 5). Among different types of fetal loss (abortion, IUFD and elective termination), elective termination of

pregnancy had the highest frequency in samples with NT ≥ 3 (33.3%). The Comparison of association of chemical parameters (PAPP-A and Free-beta HCG) and high NT with different types of fetal loss showed high frequency of abortion, IUFD and elective termination of pregnancy in high NT fetuses in comparison to low PAPP-A and Free β-HCG fetuses (Figure 1). Because the heart defect had the highest frequency in fetus with NT ≥ 3, we checked distinct subtypes of heart defects in the mentioned samples. Our results showed that frequency of ventricular septal defect (VSD) was higher than other defects (Table 6).

Discussion

In this prospective study, the results of comparison of maternal serum biochemical markers and ultrasound markers of first-trimester screening for pregnancy outcomes were evaluated for 1500 Iranian pregnant women. 2.8% of fetuses had chromosomal abnormality, specially trisomy 21 or Down syndrome. Among the mentioned screening parameters for trisomy 21, high NT had the highest detection rate (66.6%).

Table 5: Frequency of pregnancy outcome in different NT thickness

NT (mm)	≤ 3	3-4	4-5	5-6	≥ 6	Total with high NT	OR (95% CI)	P-value
	n (%)	n (%)	n (%)	n (%)	n (%)			
Successful delivery	1305 (94)	28 (53.8)	5 (15.1)	0 (0)	0 (0)	33 (31.4)	4.12 (0.35-5.1)	< 0001
Abortion	45 (3.2)	10 (19.2)	10 (30.3)	2 (13.3)	0 (0)	22 (20.9)	3.34 (0.15-3.8)	< 0001
IUFD	19 (1.36)	5 (9.6)	7 (21.2)	3 (20)	0 (0)	15 (14.2)	1.7 (1.35-4.1)	< 0001
Termination	26 (1.86)	9 (17.3)	11(33.3)	10 (66.6)	5 (100)	35 (33.3)	5.12 (0.22-3.56)	< 0001
Total	1395	52	33	15	5	105	-	-

P-value shows significant difference in some pregnancy complication between fetus with nt≤3 and nt>3 (Hardy-Weinberg equilibrium test), IUFD: Intra uterine fetal death.

Table 6: Frequency of different heart defects in different NT thickness

NT (mm)	≤ 3	3-4	4-5	5-6	≥ 6	Total
	n (%)					
Frequency of patients with heart defect	3(0.2)	5(0.3)	3(0.2)	1(0.06)	0 (0)	12(0.8)
VSD	2(0.13)	3(0.2)	1(0.06)	0 (0)	0 (0)	6 (0.4)
VSD, ASD and aortic coarctation	1(0.06)	1(0.06)	0 (0)	0 (0)	0 (0)	2 (0.13)
VSD, ASD and tricuspid valve anomaly	1(0.06)	1(0.06)	0 (0)	0 (0)	1(0.06)	3 (0.2)
Hypo plastic left ventricle	0 (0)	0 (0)	0 (0)	0 (0)	1	1 (8.3)

Hardy–Weinberg equilibrium test. Nt: Nuchal translucency, VSD: Ventricular septal defect, ASD: Atrial septal defect.




Figure 1: (a): Shows comparison of frequency of abortion in papp-a ≤ 0.4, free beta hcg ≤ 0.5 and NT > 3. (b): shows comparison of frequency of IUFD in papp-a ≤ 0.4, free beta hcg ≤ 0.5 and NT > 3. (c): shows comparison of frequency of elective termination of pregnancy in papp-a ≤ 0.4, free beta hcg ≤ 0.5 and NT > 3.

Based on our results, low level of Papp-A and free β-hCG had the most adverse effect on pregnancy outcomes in comparison to high levels of them. In this regard another multicenter study has shown that in women with high PAPP-A and free β-hCG pregnancy outcomes do not differ with those with normal levels (15). But when we analyzed the effect of low PAPP-A and free β-hCG levels on pregnancy complications, we observed impressive association of low PAPP-A on the IUGR and pre-term labor in comparison to low levels of free β-hCG. Generally speaking, PAPP-A is one of the chorionic products and is secreted in maternal blood during pregnancy. The rate of this protein grows in maternal serum during pregnancy (16, 17). During the first trimester, the decrease of this protein in mothers’ serum is one of the evidences of chromosomal aberration (18). But the predicting value of PAPP-A for pregnancy outcomes has not been reported (19).

Based on our results, the highest association of low level PAPP-A was seen with IUGR. In this context, other reports have shown IUGR to be more associated with low maternal serum PAPP-A (20, 21). Only in one research no association between low level PAPP-A and IUGR is reported (22). Concerning IUFD it can be said that this phenomenon occurs one in 160 pregnancies in developed countries. The correlation of other factors such as obesity, age, smoking, diabetes, hypertension, etc. with the IUFD was already reported, but in some studies it is showed that PAPP-A ≤ 0.4 is associated with stillbirth and IUFD. In 2002, among the 8839 pregnant women, Smith et al. reported that women with first and early second trimester PAPP-A levels in the lowest 5th centile were more likely to experience intrauterine death (23).

In another study, Dugoff et al. in 2004, concluded that first trimester low PAPP-A levels are associated with intrauterine fetal death at ≤ 24 weeks of gestation (24). Kajomaa et al. in 2017, reported the same results (25). However, the notable finding of this study was high frequency of fetal loss in fetuses with NT ≥ 3 in comparison to low levels of

PAPP-A and free-beta HCG. In this regard, we saw significant high frequency of abortion, IUFD and elective termination in fetuses with $NT \geq 3$ in comparison to low PAPP-A and free β -hCG. In this context, Dugoff et al., 2004 showed significant association of high NT with abortion in comparison to low free β -hCG among the USA pregnant women (24). In the same year among the USA pregnant women, Goetz et al. reported the association of low PAPP-A, free β -hCG and high NT with abortion but they did not compare the association of these markers individually (1). Besides, Lithner et al., 2016 reported association of high NT with miscarriage in Sweden population (26).

In our study, evaluating the association of $NT \geq 3$ with pregnancy outcomes showed that there was high frequency of elective termination of pregnancy, some of them was because of structural abnormalities such as diaphragmatic hernia, exomphalos, skeletal defects, and some genetic syndromes. Among the structural abnormalities, heart defects especially VSD had the most frequency.

Frequency of structural abnormality in our studied fetuses was 9.8% which was higher than expected in general population (2-3%) (27). Heart defects were confirmed in 12 out of 20 infants with structural abnormalities. In this regard different studies reported association of NT thickness with major heart defects when compared to those with normal hearts (28-30). For example, in a retrospective study it was shown that the 55% of heart and vessel defects were associated with increased NT (31). For these reasons, fetuses with increased NT and normal karyotype are candidate of fetal echocardiography.

Conclusion

In conclusion, based on the results of this research, among the three first trimester screening markers, high NT had the most association with different types of fetal loss and fetus structural abnormality. Regarding the fact that the first trimester screening markers have association with pregnancy complications, it is suggested that perinatologists consider this problem and when interpreting the screening results pay attention to the values of biomarkers MOM in addition to analyzing the risk of chromosomal syndromes.

Conflict of Interests

Authors have no conflict of interests.

Acknowledgments

The authors wish to express their sincere thanks to all patients and normal individuals for their contribution.

References

1. Goetzl L, Krantz D, Simpson JL, Silver RK, Zachary JM, Pergament E, et al. Pregnancy-associated plasma protein A, free β -hCG, nuchal translucency, and risk of pregnancy loss. *Obstet Gynecol* 2004; 104:30-6.
2. Schuchter K, Hafner E, Stangl G, Metznerbauer M, Ho finger D, Philipp K. The first trimester 'combined test' for the detection of Down syndrome pregnancies in 4939 unselected pregnancies. *Prenat Diagn* 2002; 22: 211-15.
3. Lawrence JB, Oxvig C, Overgaard MT, Sottrup-jensen L, GleicGJ, Hays LG, et al. The insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease secreted by human fibroblasts is pregnancy-associated plasma protein-A. *Proc Natl Acad Sci USA*. 1999; 96: 3149-53.
4. Kalousova M, Muravska A, Zima T. Pregnancy-associated plasma protein A (PAPP-A) and preeclampsia. *Adv Clin Chem* 2014; 63:169-209.
5. Smith GC, Shah I, Crossley JA, Aitken DA, Pell JP, Nelson SM, et al. Pregnancy-associated plasma protein A and alpha-fetoprotein and prediction of adverse perinatal outcome. *Obstet Gynecol* 2006; 107: 161-6.
6. Tonskaker T, Bartlett G, Trpkow C. Health information on the Internet. Gold mine or minefield? *Can FAM Physician* 2014; 60: 407-8.
7. Butler SA, Iles RK. The free monomeric beta subunit of human chorionic gonadotrophin (hCG beta) and the recently identified homodimeric beta-beta subunit (hCG beta beta) both have autocrine growth effects. *Tumour Biol* 2004; 25:18-23.
8. Ulf-Hakan S, Aila T, Henrik A, Leena V. The classification, functions and clinical use of different isoforms of hCG. *Hum Reprod Update* 2006; 12: 769-84.
9. Cruz J, Cruz G, Minerikawa R, Mazi N, Nicolaidis KH. Effect of temperature on free b-human chorionic gonadotropin and pregnancy-associated plasma protein-A concentration. *Ultrasound Obstet Gynecol* 2010; 36: 141-6.
10. Bogers H, Rifouna MS, Cohen-Overbeek TE, Koning AHJ, Willemsen SP, van der Spek PJ, et al. First trimester physiological development of the fetal foot position using three-dimensional ultrasound in virtual reality. *J Obstet Gynaecol Res* 2019; 45: 280-8.
11. Liu SS, Lee FK, Lee JL, Tsai MS, Cheng ML, She BQ, Chen SC. Pregnancy outcomes in unselected singleton pregnant women with an increased risk of first-

- trimester Down's syndrome. *Acta Obstet Gynecol Scand* 2004; 83: 1130–4.
12. Rozenberg P, Bussi eres L, Chevret S, Bernard JP, Malagrida L, Cuckle H, et al. Screening for Down syndrome using first-trimester combined screening followed by second-trimester ultrasound examination in an unselected population. *Am J Obstet Gynecol* 2006; 195: 1379–87.
 13. Snijders RJ, Noble P, Sebire N, Souka A, Nicolades KH. UK multicentre project on assessment of risk for trisomy 21 by maternal age and fetal nuchal translucency thickness at 10 – 14 weeks of gestation. *Lancet* 1998; 352: 343-6.
 14. Rocha ADS, Bernardi JR, Matos S, Kretzer DC, Sch offel AC, Goldani MZ, et al. Maternal visceral adipose tissue during the first half of pregnancy predicts gestational diabetes at the time of delivery - a cohort study. *PLoS One* 2020; 15: e0232155.
 15. Cuckle H, Arbuzova S, Spencer K, Crossley J, Barkai G, Krantz D, et al. Frequency and clinical consequences of extremely high maternal serum PAPP-A levels. *Prenat Diagn* 2003; 23: 385-8.
 16. Conover CA, Bale LK, Overgaard MT, Johnstone EW, Laursen UH, Fuchtbauer EM, et al. Metalloproteinase pregnancy-associated plasma protein A is a critical growth regulatory factor during fetal development. *Development* 2004; 131:1187-94.
 17. Lawrence JB, Oxvig C, Overgaard MT, Sottrup-Jensen L, Gleich GJ, Hays LG, et al. The insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease secreted by human fibroblasts is pregnancy-associated plasma protein-A. *Proc Natl Acad Sci USA* 1999; 96: 3149-53.
 18. Huynh L, Kingdom J, Akhtar S. Low pregnancy-associated plasma protein A level in the first trimester. *Can Fam Physician* 2014; 60:899-903.
 19. Saruhan Z, Ozakinci M, Simsek M, Mendilcioglu I. Association of first trimester low PAPP-A with adverse pregnancy outcomes. *Clin Exp Obstet Gynecol* 2012; 39: 225-8.
 20. Shah KH, Anjum A, Nair P, Bhat P, Bhat RG, Bhat S. Pregnancy associated plasma protein A: An indicator of adverse obstetric outcomes in a South India population. *Turk J Obstet Gynecol* 2020; 17: 40-5.
 21. Tul N, Pusenjak S, Osredkar J, Spencer K, Novak-Antolic Z. Predicting complications of pregnancy with first-trimester maternal serum free-betaHCG, PAPP-A and inhibin-A. *Prenat Diagn* 2003; 23: 990-6.
 22. Mendoza M, Garcia-Manau P, Ar evalo S, Avil es M, Serrano B, S anchez-Dur an M A. Diagnostic accuracy of first-trimester combined screening for early-onset and preterm preeclampsia at 8-10 weeks compared to 11-13 weeks gestation. *Ultrasound Obstet Gynecol* 2020.
 23. Smith GCS, Stenhouse EJ, Crossley JA, Aitken DA, Cameron AD, Connor JM. Early pregnancy levels of pregnancy-associated plasma protein a and the risk of intrauterine growth restriction, premature birth, preeclampsia, and stillbirth. *J Clin Endocrinol Metab* 2002; 87: 1762-7.
 24. Dugoff L, Hobbins JC, Malone FD, Porter TF, Luthy D, Comstock CH, et al. First-trimester maternal serum PAPP-A and free-beta subunit human chorionic gonadotropin concentrations and nuchal translucency are associated with obstetric complications: a population-based screening study (the FASTER Trial). *Am J Obstet Gynecol* 2004; 191: 1446-51.
 25. Kaijomaa M, Rahkonen L, Ulander VM, H am al ainen E, Alftan H, Markkanen H, et al. Low maternal pregnancy-associated plasma protein A during the first trimester of pregnancy and pregnancy outcomes. *Int J Gynaecol Obstet* 2017; 136: 76-82.
 26. Lithner CU, Kublickas M, Ek S. Pregnancy outcome for fetuses with increased nuchal translucency but normal karyotype. *J Med Screen* 2016; 23:1-6.
 27. Westin M, Saltvedt S, Bergman G, Almstrom H, Grunewlad C, Valentin L. Is measurement of nuchal translucency thickness a useful screening tool for heart defects? A study of 16,383 fetuses. *Ultrasound Obstet Gynecol* 2006; 27: 632–9.
 28. M uller MA, Bleker OP, Bonsel GJ, Bilardo CM. Nuchal translucency screening and anxiety levels in pregnancy and puerperium. *Ultrasound Obstet Gynecol* 2006; 27: 357-61.
 29. Vogel M, Sharland GK, McElhinney DB, Zidere V, Simpson JM, Miller OI, et al. Prevalence of increased nuchal translucency in fetuses with congenital cardiac disease and a normal karyotype. *Cardiol Young* 2009; 19: 441-5.
 30. Clur SA, Ottenkamp J, Bilardo CM. The nuchal translucency and the fetal heart: a literature review. *Prenat Diagn* 2009; 29: 739-48.
 31. De Domenico R, Faraci M, Hyseni E, Di Prima FA, Valenti O, Monte S, et al. Increased nuchal translucency in normal karyotype fetuses. *J Prenat Med* 2011; 5: 23–6.

Citation: Abotorabi S, Moeini N, Moghbelinejad S. **High Frequency of Fetal Loss in Fetuses With Normal Karyotype and Nuchal Translucency \geq 3 Among the Iranian Pregnant Women.** *J Fam Reprod Health* 2020; 14(2): 81-7.