

Increased Estradiol Levels Effect on Leptin Expression in Midluteal Endometrial Tissue of *Macaca Nemestrina* (Southern Pig-Tailed Macaques)

Nurhuda Sahar; Ph.D.¹, Adriana Viola Miranda; B.Med.², Afif Rasyad; B.Med.², Karina Rahمانingrum; B.Med.², Kusmardi Kusmardi; Ph.D.³, Diah Kristanty; M.Biomed.⁴

1 Department of Biology, Faculty of Medicine, University of Indonesia, Jakarta, Indonesia

2 Faculty of Medicine, University of Indonesia, Jakarta, Indonesia

3 Department of Anatomical Pathology, Faculty of Medicine, University of Indonesia, Jakarta, Indonesia

4 Department of Clinical Pathology, Faculty of Medicine, University of Indonesia, Jakarta, Indonesia

Received August 2020; Revised and accepted November 2020

Abstract

Objective: To measure the effect of serum estradiol (E2) levels on leptin expression in *Macaca nemestrina* (southern pig-tailed macaque) endometrial tissue.

Materials and methods: This study used paraffin-embedded midluteal phase endometrial tissue blocks of *Macaca nemestrina* from previous study. Included subjects were 15 female macaques of reproductive age (8-10 years) with a previous history of producing offspring, which were divided into four groups: groups administered with 30 IU, 50 IU, and 70 IU r-FSH (intervention group), and no r-FSH (control group). The stimulation was done following GnRH agonist long protocol. Staining was done using immunohistochemistry. Leptin expression was measured using immunohistochemistry (IHC) Profiler plugin of ImageJ software and counted semi-quantitatively as Histological Score (Hscore).

Results: Correlation between E2 concentration to stromal leptin expression was observed ($p=0.043$).

Conclusion: Serum estradiol concentration is found to be correlated with leptin expression in *Macaca nemestrina*, suggesting a mechanism of decreasing endometrial receptivity among women undergoing controlled ovarian hyperstimulation.

Keywords: In Vitro Fertilization; Controlled Ovarian Hyperstimulation; Leptin; Endometrial Receptivity

Introduction

Controlled ovarian hyperstimulation (COH) is known to play a crucial role in In Vitro Fertilization and Embryo Transfer (IVF-ET) due to its ability to provide a high number of viable oocytes, thus

compensating biological limitations and laboratory errors during further IVF-ET procedure (1). However, its effect on the hypothalamus-pituitary-ovarian (HPO) axis is adversely associated with endometrial receptivity (2). Several clinical studies suggest that supraphysiological levels of estradiol (E2) during midluteal phase are detrimental to embryonic implantation (3). Despite this knowledge, the causation of this effect remains controversial.

Correspondence:

Adriana Viola Miranda

Email: adriana.viola@ui.ac.id

Copyright © 2020 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-Noncommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

One of the suggested mechanisms is the role of E2 in increasing leptin expression. Leptin is an adipokine hormone initially considered to regulate food intake and energy balance (4), but experimental evidence has shown that it is also crucial in various metabolic and endocrine processes, including reproductive function. Leptin negatively affects endometrial receptivity through its inhibitory role during the thickening of the endometrial lining, uterine fluid removal, and decidualization and maternal-fetal communication (5-7). Despite this knowledge, studies on estradiol regulation of leptin expression in endometrial tissue remain limited. This study aimed to observe the correlation between leptin expression levels in the midluteal phase endometrial tissue of *Macaca nemestrina* (southern pig-tailed macaques) and estradiol concentration.

Materials and methods

Subject characteristics: This is an experimental study. We used paraffin-embedded tissue blocks of a previous study involving the uterus of *Macaca nemestrina* (southern pig-tailed macaques). The species has a similar cycle and hormonal levels with humans: their cycle length and luteal phase length are approximately one-month and 15 days, respectively (8). Subjects included in this study are 16 female macaques of reproductive age (8-10 years, 10-15 kg) with a history of producing offspring. They were tattooed with an identification number in the groin area, housed in individual cages of stainless material in a dedicated room. All animals were quarantined and adapted to new individual cages for two to three menstrual periods. During this time, animal health was maintained and any treatment was administered as needed. In the final evaluation, one subject was excluded since its menstrual cycle failed to stabilize after 3 stimulation cycles.

The animals were obtained from the Primates Animal Study Center of the Bogor Agricultural Institute, Bogor, Indonesia. All procedures involving animals performed in the study were approved by the Institutional Animal Care and Use Committee for Primate Animal Studies of Bogor Agricultural Institute from a previous study (ACUC no. 08-B001-1R, 1 February 2010) which is further stated in Letter No.685/IT3.L1.9/TU/2019, 24 October 2019. This research was part of a bigger project where the majority of the organs of *Macaca nemestrina* were used.

These macaques were divided into four groups in accordance with their COH protocols, which include

administration of (Gonadotropin-releasing hormone) GnRH agonist and recombinant Follicle-stimulating hormone (FSH) (r-FSH) with dosages of 30 IU, 50 IU, 70 IU (intervention groups), and no r-FSH (control group).

Controlled ovarian hyperstimulation protocol: In accordance with GnRH agonist long protocol, 160 µg GnRH agonist (Suprefacts; Sanofi SA Paris, France) was administered for approximately 14 days, from the beginning of the luteal phase of the previous menstrual cycle until the day before ovulation. On the second day after menstruation, if the estradiol level reached <70 pg/mL, recombinant FSH (Gonal-F; Merck KGa Darmstadt, Germany) was injected with three different doses according to the treatment groups (30, 50, and 70 IU) per day for 10 or 12 days until estradiol secretion's peak. Then, to stimulate ovulation, HCG (pregnyl; Merck KGa) was administered at a dosage of 3200 IU (equivalent to 10,000 IU for humans). The luteal phase was determined by measuring serial progesterone levels starting on post-ovulation day. Endometrial tissues were then retrieved on day 20-22 of the midluteal phase and embedded in paraffin blocks.

Sample examination: The staining of endometrial tissues retrieved from paraffin blocks was conducted using immunohistochemistry method. Firstly, endometrial tissues of *Macaca nemestrina* embedded in paraffin blocks were cut with a thickness of 3,5 µm. Antigen retrieval was done in a container containing Tris Ethylenediaminetetraacetic acid (EDTA) (pH=9) which was heated in Retrieval Generation One BioGear with a temperature of 98°C for 15 minutes. Incubation with leptin monoclonal antibody was conducted in a 4°C temperature for 24 hours (overnight). E2 serum concentrations were obtained from our previous study (9).

Image acquisition: For each tissue, we measured the immunoreactivity of three endometrial components: glandular, stromal, and luminal components. Images were obtained using a camera and IndoMicroView software. Images were taken using x40 objective, with x1, 5 magnifications.

Data extraction: Endometrial component images were exported from IndoMicroView software as TIF images. Immunoreactivity to leptin is divided into four groups based on the number of positive cells: negative, low positive, positive, and high positive. The percentage of each cell group was quantified automatically with IHC Profiler plugin of ImageJ. Each component immunoreactivity was measured on

four different slide areas, randomly picked by the authors. These percentages were used to score the histological score (Hscore) for each slide, with a formula as follows (pi is the percentage of cells with i stain, with i being immunoreactivity).

$$\text{Histological Score} = \sum p_i \times i$$

Statistical analysis: Data were presented as mean. Comparison of Hscore between control and intervention groups of each compartment was analyzed parametrically with Analysis of variance (ANOVA) and nonparametrically with the Kruskal-Wallis test, using SPSS release 22. Correlation between groups of E2 concentration and leptin expression was further analyzed with the Kruskal-Wallis test. Statistical decision was done using 5% significance level (p=0.05).

Results

Immunolocalization of leptin expression in *Macaca nemestrina* endometrium

The expression of leptin in the mid-luteal endometrium phase of *Macaca nemestrina* was found in the cytoplasm of glandular epithelial cells, stromal cells, and luminal epithelial cells, with the level being consistently higher in the latter (Figure 1). These results were in line with leptin expression in the endometrium with natural cycle (6).




Figure 1: Immunohistochemical localization of leptin expression in *Macaca nemestrina* endometrium. Localization of leptin expression in cytoplasmic glandular epithelium cells (A), stromal cells (B), and luminal epithelial cells (C) in 400x magnification. Positive control (D) and negative control (E) are also shown.

Leptin expression between different stimulation protocols

Our result found no significant differences in leptin expression in the endometrial tissue between different stimulation protocols (Figure 2).




Figure 2: Graphical representation of leptin expression (quantified as Hscore). No significant differences in leptin expression were found between different stimulation protocols.

To illustrate the effect of estradiol levels on endometrial leptin expression, based on receiver operating characteristic (ROC) curve analysis on the leptin and estradiol data, we divided the concentration levels into two groups: E2<1000 pg/mL and E2≥1000 pg/mL. We observed a correlation between E2 concentration to stromal leptin expression (p=0.043). No significant differences were found in glandular (p=0.70) and luminal epithelial (p=0.405) leptin expression (Figure 3).

Discussion

Leptin is known to be crucial in determining endometrial receptivity. Several leptin systems associated with this association have been proposed, including γ -Epithelial sodium channel (γ -ENaC) regulation in luminal epithelial cells (5), endometrial thickness alteration through leptin activities in luminal and glandular epithelia (10), and decidualization and mother-embryo communication inhibition in stromal components of the tissue (11). Therefore, alterations of endometrial leptin expression levels after controlled ovarian hyperstimulation (COH) are postulated to cause decreased endometrial receptivity and thus, IVF success rate.

Several studies have shown evidence for the causation. Higher levels of leptin in serum (12) and follicular fluid (6, 7, 13) are found to reduce

pregnancy rate within women undergoing IVF. Our findings, however, observe no significant differences in endometrial leptin expression between different COH protocols.




Figure 3. Average leptin expression within three E2 concentration groups: control, E2<1000, and E2>1000 in three endometrial tissue components: (a) glandular component; (b) stromal component; and (c) luminal epithelial component. Significant differences in leptin expression are shown in the stromal component.

The regulator of leptin expression in endometrium

remains somewhat controversial. Several studies show that estradiol increases serum leptin levels within normal menstrual cycles. Another study on human endometrial tissue within different menstrual phases also supports this conclusion (14). Several studies, contradict these findings: no significant correlations was seen between serum levels of both hormones during IVF cycles, and hormone replacement therapy (HRT) of estradiol in postmenopausal women with leptin concentration (13, 15). The result of this study shows that leptin expression is correlated with higher E2 concentration (>1000 pg/mL), thus it can be postulated that estradiol acts upon endometrial leptin expression, which then exerts its effect on endometrial receptivity. In a previous study, estradiol was also found to be correlated with the expression of leptin receptor (Ob-R), suggesting its large influence on the leptin system in the reproductive organs (16).

Our study has several limitations: small sample size, limitation on standardizing the duration of each menstrual cycle phase. We also did not measure the ovarian reserve of the subjects. These limitations may affect the result of our study since they are known to affect the hormonal profile of endometrial tissues.

Conclusion

Serum estradiol concentration is found to be correlated with leptin expression in *Macaca nemestrina*, suggesting a mechanism of decreasing endometrial receptivity among women undergoing controlled ovarian hyperstimulation.

Conflict of Interests

Authors have no conflict of interests.

Acknowledgments

We would like to thank the Directorate of Research and Community Engagement (DRPM) Universitas Indonesia for the PITTA B 2019 research grant.

References

1. Heger A, Sator M, Pietrowski D. Endometrial receptivity and its predictive value for IVF/ICSI-outcome. *Geburtshilfe Frauenheilkd* 2012; 72: 710–5.
2. Dodson WC, Haney AF. Controlled ovarian hyperstimulation and intrauterine insemination for treatment of infertility. *Fertil Steril* 1991; 55: 457–67.
3. Joo BS, Park SH, An BM, Kim KS, Moon SE, Moon HS. Serum estradiol levels during controlled ovarian

- hyperstimulation influence the pregnancy outcome of in vitro fertilization in a concentration-dependent manner. *Fertil Steril* 2010; 93: 442–6.
4. Catteau A, Caillon H, Barrière P, Denis MG, Masson D, Fréour T. Leptin and its potential interest in assisted reproduction cycles. *Hum Reprod Update* 2016; 22: 320-41.
 5. Lin X-H, Liu M-E, Xu H-Y, Chen X-J, Wang H, Tian S, et al. Leptin down-regulates γ -ENaC expression: a novel mechanism involved in low endometrial receptivity. *Fertil Steril* 2015; 103: 228-235.e3.
 6. Chakrabarti J, Chatterjee R, Goswami S, Chakravarty B, Kabir SN. Overt leptin response to controlled ovarian hyperstimulation negatively correlates with pregnancy outcome in in vitro fertilization-embryo transfer cycle. *J Hum Reprod Sci* 2012; 5: 194–9.
 7. Hestiantoro A, Wiweko B, Purwaka RH. Basal follicle stimulating hormone and leptin on the day of hCG administration predict successful fertilization in in vitro fertilization. *Med J Indones* 2016; 25: 39.
 8. Blakley GB, Beamer TW, Dukelow WR. Characteristics of the menstrual cycle in nonhuman primates. IV. Timed mating in *Macaca nemestrina*. *Lab Anim* 1981; 15: 351–3.
 9. Sahar N, Mujihartini N, Pudjianto DA, Pradhita AD, Thu R, Kusmardi K. Increased progesterone on the day of administration of hCG in controlled ovarian hyperstimulation affects the expression of HOXA10 in primates' endometrial receptivity. *Biomedicines* 2019; 7(4):83.
 10. Chou SH, Mantzoros C. 20 Years of Leptin: Role of leptin in human reproductive disorders. *J Endocrinol* 2014; 223: T49-62.
 11. Tanaka T, Utsunomiya T, Bai T, Nakajima S, Umesaki N. Leptin inhibits decidualization and enhances cell viability of normal human endometrial stromal cells. *Int J Mol Med* 2003; 12: 95–8.
 12. Kaplanoğlu İ, Kaplanoğlu GT, Çınar Ö. Is leptin receptor expression triggered in the case of embryo transfer to endometrium coculture? *Turk J Med Sci* 2019; 49: 1014–8.
 13. Geber S, Brandão AHF, Sampaio M. Effects of estradiol and FSH on leptin levels in women with suppressed pituitary. *Reprod Biol Endocrinol* 2012; 10: 45.
 14. Özler A, Kuşçu NK, Temiz P, Kandiloğlu AR, Koyuncu FM. Leptin expression in proliferative, secretory and hyperplastic endometrial tissues. *J Turk Ger Gynecol Assoc* 2011; 12: 157–61.
 15. Hill MJ, Uyehara CFT, Hashiro GM, Frattarelli JL. The utility of serum leptin and follicular fluid leptin, estradiol, and progesterone levels during an in vitro fertilization cycle. *J Assist Reprod Genet* 2007; 24: 183-8.
 16. Méndez-López L, Zavala-Pompa A, Fuentes-Vera M, Pacheco-Calleros J, Escobar-Moreno A, Cortés-Gutiérrez E, et al. Quantification of leptin receptor expression in endometrial tissue throughout the human menstrual cycle. *Gen Endocrinol* 2015; 11: 147–54.

Citation: Sahar N, Miranda AV, Rasyad A, Rahmaningrum K, Kusmardi K, Kristanty D. **Increased Estradiol Levels Effect on Leptin Expression in Midluteal Endometrial Tissue of *Macaca Nemestrina* (Southern Pig-Tailed Macaques).** *J Fam Reprod Health* 2020; 14(4): 229-33.