

# Maternal Sleep and Related Pregnancy Outcomes: A Multicenter Cross-Sectional Study in 11 Provinces of Iran

Mahmoud Hajipour; M.Sc.<sup>1</sup>, Maryam Soltani; M.Sc.<sup>2</sup>, Roya Safari-Faramani; Ph.D.<sup>3</sup>,  
Salman Khazaei; Ph.D.<sup>4</sup>, Koorosh Etemad; Ph.D.<sup>5</sup>, Sharmin Rahmani; M.Sc.<sup>6</sup>,  
Tannaz Valadbeigi; M.Sc.<sup>7</sup>, Halime Yaghoobi; M.Sc.<sup>8</sup>, Shahab Rezaeian; Ph.D.<sup>9,10</sup>

1 Research Center Office, Epidemiology Department, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2 Razi Clinical Research Development Unit (RCRDU), Birjand University of Medical Sciences (BUMS), Birjand, Iran

3 Research Center for Environmental Determinants of Health, Health Institute, School of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

4 Research Center for Health Sciences, Hamadan University of Medical Sciences, Hamadan, Iran

5 Department of Epidemiology, Environmental and Occupational Hazards Control Research Center, Faculty of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

6 Student Research Committee, Research Center for Environmental Determinants of Health, School of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

7 Clinical Research Development Unit, Imam Hossein Hospital, Shahroud University of Medical Sciences, Shahroud, Iran

8 Social Determinants in Health Promotion Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

9 Infectious Diseases Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

10 Clinical Research Development Center, Imam Reza Hospital, Kermanshah University of Medical Sciences, Kermanshah, Iran

Received November 2020; Revised and accepted December 2020

## Abstract

**Objective:** Sleep disturbance during pregnancy is one of the most common maternal complaints. Not only does it play a crucial role in a mother's life, but also it comes with a multitude number of complications. This study aimed at assessing the association between sleep disturbance in pregnancy and maternal and child outcomes.

**Materials and methods:** This was a multicenter cross-sectional study, conducted on pregnant women across 11 provinces in Iran in 2018. Sleep disturbance as a composite variable was defined using the principal component analysis based on five questions. Abortion, anemia in the first and third trimester, gestational diabetes, gestational age, glucose tolerance test (GTT), fasting blood sugar (FBS), mode of delivery, low birth weight and stillbirth were defined as study outcomes.

**Results:** Totally, 3675 pregnant women enrolled in the study. Most of the participants (84.5%) reported

## Correspondence:

Dr. Shahab Rezaeian

Email: shahab.rezaeian@kums.ac.ir

Copyright © 2021 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-Noncommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

that their sleep duration is less than 8 hours per day. The prevalence of sleep disturbance was 20.7% (95% CI: 19.1, 22.3). After adjusting for maternal age, education, job, place of residency and physical violence, sleep disturbance would increase the odds of abortion ( $p=0.009$ ), anemia in both first ( $p=0.001$ ) and third ( $p=0.003$ ) trimester, gestational age ( $p=0.049$ ), abnormal FBS ( $p=0.015$ ) and cesarean section ( $p<0.001$ ).

**Conclusion:** Regarding the effect of sleep quality on maternal outcomes, planning and implementing a suitable intervention in the context of primary health care is necessary. Increasing the awareness of mothers, health workers and medical personnel about the suitable quality and quantity of sleep during pregnancy is of great importance.

**Keywords:** Sleep Wake Disorders; Pregnancy Outcome; Stillbirth; Iran

## Introduction

Sleep is a bio-behavioral process that is essential for the human body to have a proper performance (1). It should be noted that sleep does not mean just the absence of waking. But also, during this process, specialized physiological activities occur in the brain and throughout the body. In fact, through this active process, tissue repair, memory stabilization, and homeostatic balance in the body are maintained (2).

Results of several studies have shown that there is a U-shape relationship between the duration of sleep and death; so that seven hours of sleep per day is associated with the lowest mortality rate (3). Evidence indicate a link between sleep disturbances and increased risk of depression, diabetes, obesity as well as all-causes of mortality (3-6).

Sleep disturbances are a common issue during pregnancy so that globally more than 75% of pregnant women experience some form of sleep problems (7). Rising levels of hormones such as progesterone can be responsible for such changes in the quality of sleep in pregnant women (8). In pregnant women, short sleep duration is associated with longer labor times and subsequently increased cesarean section (C-section) rates (9). Along with sleep disorders in pregnant women, risen risk of some complications such as preeclampsia, hypertension, and small for gestational age (SGA) infants have been reported (10, 11). Besides, the quality of life in pregnant women is related to the quality of sleep (12).

Identifying risk factors of developing maternal and fetal adverse outcomes in pregnant women is essential for preventive strategies. There is limited evidence regarding maternal and fetal outcomes in pregnant women with sleep disturbances in Iran. This study was conducted to assess the relationship between sleep disturbance in pregnancy and related

pregnancy outcomes in Iran.

## Materials and methods

**Study design and setting:** This was a multicenter cross-sectional study, a part of case control study in 2018 (13), which designed to determine the relationship between sleep disturbance and adverse pregnancy outcomes among mothers referring to the health centers in eleven provinces/cities across Iran including Fars, Hormozgan, Kermanshah, Chaharmahal and Bakhtiari, Hamadan, Kohgiluyeh and Boyer-Ahmad, Yazd, South Khorasan and Golestan provinces and two big cities of Mashhad and Zahedan.

**Study population and sampling procedure:** In this study, we have tried to sample from different regions of Iran, so we have chosen different provinces for study to enable the generalization of the published results to the entire Iranian population. To this end, all the provinces were divided into four different geographical regions (North, South, East, and West). Then, two health centers as one urban and one rural were selected from each geographical region from two big cities. Lastly, 20 subjects were randomly selected from each health center. Some data, mentioned in the data collection section, were collected by interviewing the mothers and some were also extracted from family health records by using a checklist. Totally, a sample size of 3675 pregnant women was enrolled in the study. Having a health record was an inclusion criterion for our study and those with missing data were excluded in the analysis. Women were invited to interview by telephone.

**Data collection methods and tools:** The data collection form in this study was a researcher-made checklist that includes the mother's personal information such as age, education, body mass index (BMI), place of residency, occupation, domestic violence, and daily physical activity.

In this study, sleep disturbance was defined by

five questions including the length of sleeping (in an hours), sleep location, having regular sleep, sleep position during the last pregnancy and self-reported sleep status as very good, good, bad and very bad. Sleep disturbance as a composite variable was defined using the principal component analysis (PCA) and was classified as dichotomous variable for analysis. In addition, length of sleeping (in hour) was categorized into two categories as lower than 8 (short sleep) and equal to or more than 8 hours.

Dependent variables (study outcomes) were gestational age (<37 or ≥37 weeks), gestational diabetes mellitus (GDM), maternal anemia in the first or third trimesters (women with hemoglobin level lower than 11 g/dL), low birth weight (lower than 2500 g), abortion, stillbirth, mode of delivery (C-section or vaginal), glucose tolerance test (GTT) and fasting blood sugar (FBS) abnormal. GDM, GTT- and FBS-abnormal in pregnancy were defined according to WHO criteria (14).

Although, based on the Centers for Disease Control and Prevention (CDC), stillbirth is classified into three categories, including early stillbirth (death at 20 to 27 weeks of pregnancy), late stillbirth (death at 28 to 36 weeks of pregnancy) and term stillbirth (death in the 37th or 38th week of pregnancy or at delivery), it was not specific in our study. The stillbirth was collected as a dichotomous variable.

**Data analysis:** The statistical analyzes used in this study were descriptive analyzes and logistic regression model at a significant level of 5%. In the adjusted models, the odds ratios were adjusted for maternal variables including maternal age, education, job, place of residency, physical activity and domestic violence.

All analyses were done by STATA software version 14 (STATA Corp, College Station, TX).

**Ethical considerations:** To obtain information on infant mortality and information on household records, after being confirmed at the University, the correspondence was presented to coordinate the universities. All information was kept confidential and was prevented by the name and surname in the reporting phase. Verbal informed consent was obtained from all mothers.

## Results

Totally, 3675 pregnant women enrolled in the study. The mean ±SD of maternal age and BMI was 27.2±6.0 and 23.1±4.8, respectively. Mothers with short sleep were significantly older ( $p<0.001$ ), higher BMI ( $p<0.005$ ) and less educated ( $p<0.001$ ). The proportion of short sleep was 20.3, 10.9 and 13.7 percent in participants with primary, secondary, high school and academic education. Short sleep was as common in urban residents as rural ones ( $p=0.123$ ). Mean ±SD of sleep time was 7.3±2.7 in participants with less than 8 hours sleep which was significantly two hours less than the other group ( $p<0.001$ ) (Table 1).

Based on the PCA method, the prevalence of sleep disturbance was 20.7% (95% CI: 19.1, 22.3). Most of the participants (84.5%) reported that their sleep duration is less than 8 hours per day. Bad and very bad sleep qualities were reported by 2.8 and 2.8 percent, respectively. Around 40 percent of the participants (38.4%) reported that they have regular sleep very often. Most of the participants reported that they prefer to sleep in dark room (42.5%) and on side position (82.2%) (Table 2).

**Table 1:** Characteristics of study participants according to sleep duration

	Sleep duration		P-value
	≥ 8 hours	< 8 hours	
Mother's age (year)	26.9±5.8	28.7±6.6	<0.001
BMI*	23.0±4.5	24.0±5.2	<0.002
Job			
Housewife	2815 (85.5)	478 (14.5)	0.089
Employed	159 (81.9)	35 (18.1)	
Self-employed	79 (79.0)	21 (21.0)	
Education			<0.001
Primary	846 (79.7)	215 (20.3)	
Secondary	803 (89.1)	98 (10.9)	
High school and academic	1428 (86.3)	226 (13.7)	
Place of residency			0.123
Rural	1661 (86.1)	269 (13.9)	
Urban	1344 (84.2)	252 (15.8)	
Sleep time (hour)	9.2±2.5	7.3±2.7	<0.001

\*Body mass index; data were presented as number (%) for categorical variables and mean±SD for continuous ones.

**Table 2:** Self-reported maternal sleep during pregnancy

Variable	No.	%	
Sleep location	Darkness	1,539	42.2
	Dark and light	1,943	53.3
	Light	125	3.4
	Missing	42	1.2
Sleep position	On side	3,000	82.2
	Supine	595	16.3
	Missing	54	1.5
Quality of sleep	Very good	322	8.8
	Good	2,760	75.6
	Bad	437	12.0
	Very Bad	102	2.8
	Missing	28	0.8
Having regular sleep	Always	465	12.7
	Very Often	1,409	38.6
	Sometimes	1,383	37.9
	Rarely	309	8.5
	Never	53	1.5
	Missing	30	0.8
Sleep duration (hour)	<8	3,082	84.5
	≥8	539	14.8
	Missing	28	0.8

After adjusting for maternal age, education, job,

place of residency, daily physical activity and domestic violence, sleep disturbance would increase the odds of abortion by 41 percent ( $p=0.009$ ).

The odds of anemia in both first ( $p=0.001$ ) and third ( $p=0.003$ ) trimesters would increase by 51 and 71 percent, respectively. Sleep disturbance also was associated with gestational age ( $p=0.049$ ) and C-section ( $p<0.001$ ).

Sleep disturbance was directly associated with abnormal FBS ( $p=0.015$ ). It also increased the odds of GGT by 35 percent ( $p=0.065$ ). Although the odds of low birth weight (LBW) would increase by 28 percent due to sleep disturbance, it did not reach to statistical significance (Table 3).

## Discussion

A prevalence of 20.7% was found for sleep disturbance in the Iranian pregnant women. We also found an important relationship between sleep disturbance during pregnancy and some adverse pregnancy outcomes such as abortion, anemia in both first and third trimesters, gestational age, C-section and abnormal FBS, after adjusting for maternal age, education, job, place of residency, daily physical activity and domestic violence.

**Table 3:** The adjusted odds ratios of the association between sleep disorder (as independent variable) and related pregnancy outcome in Iran

Variable	Adjusted OR (95% CI)*	P-value
Abortion	No	Ref.
	Yes	1.41 (1.09, 1.82)
Anemia in the 3 <sup>rd</sup> trimester	No	Ref.
	Yes	1.51 (1.19, 1.92)
Anemia in the 1 <sup>st</sup> trimester	No	Ref.
	Yes	0.71 (0.57, 0.89)
Gestational age	≥37 weeks	Ref.
	<37 weeks	1.31 (1, 1.7)
Gestational diabetes	No	Ref.
	Yes	1.24 (0.8, 1.93)
GTT	Normal	Ref.
	Abnormal	1.37 (0.98, 1.92)
FBS	Normal	Ref.
	Abnormal	1.54 (1.09, 2.18)
Birth weight (gr)	2500-4000	Ref.
	<2500	1.28 (0.99, 1.66)
Mode of delivery	Vaginal	Ref.
	Cesarean	1.59 (1.3, 1.94)
Birth outcome	Live birth	Ref.
	Stillbirth	0.94 (0.74, 1.19)

OR: odds ratio; GTT: glucose tolerance test; FBS: fasting blood sugar

\*In the adjusted models maternal variables including age, education, job, place of residency, physical activity and domestic violence were adjusted.

Short sleep duration and poor sleep quality are common problems during pregnancy (15). This study was carried out to examine the relationship between poor sleep quality and related pregnancy outcomes. Findings of this study showed that sleep disturbance is associated with preterm delivery which is in accordance with previous studies (16-18). Okun and colleagues found that the level of interleukin-6 (IL-6) was higher in pregnant women with short sleep duration or poor sleep quality (19). They argued that higher level of Cytokines such as IL-6 was related to the initiation of spontaneous pain associated with preterm delivery (19). A number of studies revealed that sleep disorders may lead to impaired natural immune mechanisms (20). Generally, changes in sleep duration vary throughout the pregnancy. As such, sleep duration is shorter compared to the whole day sleep duration. Maternal sleep disruption is commonly due to increased levels of progesterone, nausea, vomiting, frequent urination and anorexia in the first trimester, increased uterus and urinary frequency, heart rate, and fetal movement in the second trimester, and physical changes and disability, trouble falling asleep, and common complaints such as back pain and itching in the third trimester (21-23).

The findings of the present study suggested that the odds of abortion are higher among mothers with sleep disturbance. Also, a sleep duration of 8hrs and lower increase the risk of abortion by four times and two times in the first and second trimester, respectively (24). Louis and colleagues stated that sleep apnea results in the onset of spontaneous delivery and subsequently separation of the placental membranes and abortion (25). Sleep deprivation especially during the first 20 weeks of pregnancy dramatically increase the inflammatory response and trophoblast invasion in the uterus (26). The fact that snoring and difficulty breathing cause sleep deprivation, it can be concluded that low sleep quality can lead to abortion (27).

Our results showed that the odds of cesarean was higher among women with sleep disturbance which was in accordance with previous studies (15, 28). Zafarghandi and colleagues found that sleep duration and the quality were correlated with the mode of delivery (29, 30). As such, pregnant women with longer sleep duration were more likely to give natural delivery. In contrast, Teong and colleagues in a prospective study stated that sleep measure is not a predictive factor for caesarean section (31). Li and colleagues argued that women with poor sleep quality

were 1.87, 5.19, and 1.82 times more likely to deliver by cesarean in the first, second and third trimester, respectively (30). However, Evans and colleagues found no significant relationship between sleep quality and duration with the mode of delivery (32). Natural vaginal delivery is a high energy expending process, thus sleep deprivations such as poor-quality sleep and sleep disturbance decrease the ability to perform a perfect labor especially in the last month of gestation (33). The variation in these results may be as a result of sociocultural differences, different measurement methods, and various implications of sleep as well as different sampling method. Furthermore, there are many mediating factors in which cesarean is performed, such as breech presentation, pelvic stenosis, multiple births, third trimester bleeding due to placenta previa, and severe placental abruption, and the umbilical cord prolapse. All these factors may influence the mode of delivery (30, 34). Therefore, further efforts are required to investigate all factors individually.

The findings of the present study showed that the risk of sleep disturbance was higher in the first and third trimester among pregnant women with anemia. Results from studies show that restless leg syndrome is a common disorder among women due to reduced level of hemoglobin and vitamin deficiencies especially during pregnancy. Evidences show that the sleep quality was significantly improved among recovered patients with restless leg syndrome (35).

The number of studies focusing on the sleep disturbance and diabetes is limited and the biological mechanisms affecting sleep disturbance such as short or long sleep duration, poor sleep quality, and gestational diabetes remain unknown. However, some studies have argued that short sleep duration increased the odds of gestational diabetes (36-38), which is consistent with our results. Facco and colleagues in a cohort study reported that after adjusting for age, BMI and race/ethnicity, short sleep duration is associated with gestational diabetes (38). A previous meta-analysis with including five studies showed that sleep disordered breathing increased the risk of gestational diabetes by two times (39). Observational and experimental studies have shown that long or short sleep duration lead to impaired insulin production and glucose metabolism. The mediating factors in this association include increased oxidative stress, increased systemic inflammation, disruption of energy homeostasis, hormone changes in pregnant

women, and physical problems and stress during labor (21, 40-43). However, physical inactivity and sleep deprivation during pregnancy has been shown to be associated with longer sleep duration and dysfunction of glucose metabolism (44).

Besides, the risk of LBW was higher in women with sleep disturbance. Basan et al found no significant relationship between the infant birth weight and sleep disturbance and poor sleep quality (45). A number of studies have shown a significant relationship between low birth weight and maternal sleep disturbance which was not in agreement with the present study (17, 46). In spite the many logical reasons, the relationship between poor sleep quality and LBW is still unclear. For instance, maternal sleep disturbance may be attributed to changes in maternal appetite leading to increased fetal weighing.

According to the findings of the present study, no significant relationship was observed between sleep disturbance and stillbirth and infant mortality. Factors that are significantly associated with sleep disturbance during pregnancy such as high maternal age, ethnicity, tobacco use, anemia, congenital anomaly, maternal medical condition, intrauterine growth restriction, and previous history of adverse pregnancy outcomes were overlapped with the risk factors for perinatal mortality (47, 48). This overlap has formed many reliable biological hypotheses, thus further researches to assess the relationship between sleep disturbance and stillbirth is needed.

The present study had an important limitation. A standard sleep-quality questionnaire was not used. However, we attempted to declare an estimate of sleep disturbance with a large sample among Iranian pregnant women which may increase the generalizability of the study findings.

## Conclusion

The findings of the current study revealed that sleep disturbance is a main risk factor for preterm birth, cesarean birth, gestational diabetes, anemia, and LBW. Therefore, awareness of health and medical personnel regarding the quality and quantity of sleep during pregnancy is of great importance and training intervention should be implemented to reduce the adverse outcomes of pregnancy associated with sleep disturbance.

## Conflict of Interests

Authors have no conflict of interests.

## Acknowledgments

The authors would like to thank the Clinical Research Development Center of Imam Reza Hospital for their wise advices. This study was approved by the Ethics Committee of Kermanshah University of Medical Sciences (code: 980808). The consent to participate was verbally obtained and this method was approved by the institutional review board (IRB) of the University.

## References

1. Okun ML, Roberts JM, Marsland AL, Hall M. How disturbed sleep may be a risk factor for adverse pregnancy outcomes a hypothesis. *Obstet Gynecol Surv* 2009; 64: 273-80.
2. Fuller PM, Gooley JJ, Saper CB. Neurobiology of the sleep-wake cycle: sleep architecture, circadian regulation, and regulatory feedback. *J Biol Rhythms* 2006; 21: 482-93.
3. García-Perdomo HA, Zapata-Copete J, Rojas-Cerón CA. Sleep duration and risk of all-cause mortality: a systematic review and meta-analysis. *Epidemiol Psychiatr Sci* 2019; 28: 578-88.
4. Grandner MA, Seixas A, Shetty S, Shenoy S. Sleep Duration and Diabetes Risk: Population Trends and Potential Mechanisms. *Curr Diab Rep* 2016; 16: 106.
5. Ouyang P, Sun W. Depression and sleep duration: findings from middle-aged and elderly people in China. *Public health* 2019; 166: 148-54.
6. Wu Y, Zhai L, Zhang D. Sleep duration and obesity among adults: a meta-analysis of prospective studies. *Sleep Med* 2014; 15: 1456-62.
7. Facco FL, Grobman WA, Kramer J, Ho KH, Zee PC. Self-reported short sleep duration and frequent snoring in pregnancy: impact on glucose metabolism. *Am J Obstet Gynecol* 2010; 203: 142. e1-5.
8. Lee J, Eklund EE, Lambert-Messerlian G, Palomaki GE, Butterfield K, Curran P, et al. Serum Progesterone Levels in Pregnant Women with Obstructive Sleep Apnea: A Case Control Study. *J Womens Health (Larchmt)* 2017; 26: 259-65.
9. Lee KA, Gay CL. Sleep in late pregnancy predicts length of labor and type of delivery. *Am J Obstet Gynecol* 2004; 191: 2041-6.
10. Perez-Chada D, Videla AJ, O'flaherty ME, Majul C, Catalini AM, Caballer CA, et al. Snoring, witnessed sleep apnoeas and pregnancy induced hypertension. *Acta Obstet Gynecol Scand* 2007; 86: 788-92.
11. Franklin KA, Holmgren PA, Jonsson F, Poromaa N, Stenlund H, Svanborg E. Snoring, pregnancy-induced

- hypertension, and growth retardation of the fetus. *Chest* 2000; 117: 137-41.
12. Rezaei E, Behboodi Moghadam Z, Saraylu K. Quality of life in pregnant women with sleep disorder. *J Family Reprod Health* 2013; 7 :87-93.
  13. Hajipour M, Tabatabaee HR, Etemad K, Ansari H, Ansari Moghadam A, Mohammadi M, et al. The Pre-Pregnancy Risk Factors of Stillbirth in Pregnant Iranian Women: A Population-Based Case-Control Study. *Health Scope* 2019; 8: e64034.
  14. World Health Organization. Diagnostic criteria and classification of hyperglycaemia first detected in pregnancy: a World Health Organization guideline. *Diabetes Res Clin Pract* 2014; 103: 341–63.
  15. Palagini L, Gemignani A, Banti S, Manconi M, Mauri M, Riemann D. Chronic sleep loss during pregnancy as a determinant of stress: impact on pregnancy outcome. *Sleep Med* 2014; 15: 853-9.
  16. Okun ML, Luther JF, Wisniewski SR, Sit D, Prairie BA, Wisner KL. Disturbed sleep, a novel risk factor for preterm birth? *J Womens Health (Larchmt)* 2012; 21: 54-60.
  17. Micheli K, Komninos I, Bagkeris E, Roumeliotaki T, Koutis A, Kogevinas M, et al. Sleep patterns in late pregnancy and risk of preterm birth and fetal growth restriction. *Epidemiology* 2011; 22: 738-44.
  18. Alihosseini E, Najar S, Haghighizadeh MH. The Relationship Between Sleep Disorders During Pregnancy and Miscarriage. *Jundishapur J Chronic Dis Care* 2017; 6: e41340.
  19. Okun ML, Hall M, Coussons-Read ME. Sleep disturbances increase interleukin-6 production during pregnancy: implications for pregnancy complications. *Reprod Sci* 2007; 14: 560-7.
  20. Okun ML, Luther JF, Wisniewski SR, Wisner KL. Disturbed sleep and inflammatory cytokines in depressed and nondepressed pregnant women: an exploratory analysis of pregnancy outcomes. *Psychosom Med* 2013; 75: 670-81.
  21. Lee KA, Zaffke ME, McEnany G. Parity and sleep patterns during and after pregnancy. *Obstet Gynecol* 2000; 95: 14-8.
  22. Hanif S. Frequency and pattern of urinary complaints among pregnant women. *J Coll Physicians Surg Pak* 2006; 16: 514-7.
  23. Okun ML, Coussons-Read ME. Sleep disruption during pregnancy: how does it influence serum cytokines? *J Reprod Immunol* 2007; 73: 158-65.
  24. Samaraweera Y, Abeysena C. Maternal sleep deprivation, sedentary lifestyle and cooking smoke: Risk factors for miscarriage: A case control study. *Aust N Z J Obstet Gynaecol* 2010; 50: 352-7.
  25. Louis JM, Auckley D, Sokol RJ, Mercer BM. Maternal and neonatal morbidities associated with obstructive sleep apnea complicating pregnancy. *Am J Obstet Gynecol* 2010; 202: 261.e1-5.
  26. Okun ML, Roberts JM, Marsland AL, Hall M. How disturbed sleep may be a risk factor for adverse pregnancy outcomes. *Obstet Gynecol Surv* 2009; 64: 273-80.
  27. Espiritu JR. Aging-related sleep changes. *Clin Geriatr Med* 2008; 24: 1-14, v.
  28. Lee KA, Gay CL. Sleep in late pregnancy predicts length of labor and type of delivery. *Am J Obstet Gynecol* 2004; 191: 2041-6.
  29. Zafarghandi N, Hadavand S, Davati A, Mohseni SM, Kimiaimoghadam F, Torkestani F. The effects of sleep quality and duration in late pregnancy on labor and fetal outcome. *J Matern Fetal Neonatal Med* 2012; 25: 535-7.
  30. Li R, Zhang J, Zhou R, Liu J, Dai Z, Liu D, et al. Sleep disturbances during pregnancy are associated with cesarean delivery and preterm birth. *J Matern Fetal Neonatal Med* 2017; 30: 733-8.
  31. Teong ACA, Diong AX, Omar SZ, Tan PC. The Impact of Self-Reported Sleep on Caesarean Delivery in Women Undergoing Induction of Labour: A Prospective Study. *Scientific Reports* 2017; 7: 12339.
  32. Evans ML, Dick MJ, Clark AS. Sleep during the week before labor: relationships to labor outcomes. *Clinical nursing research* 1995; 4: 238-49.
  33. Beebe KR, Lee KA. Sleep disturbance in late pregnancy and early labor. *J Perinat Neonatal Nurs* 2007; 21: 103-8.
  34. Nazarpour S. Sleep Disorders in Pregnancy and Its Effect on the Length of the Labor and Type of Delivery in Firuzabadi Hospital in Tehran in 2006. *Iran Journal of Nursing* 2007; 20: 63-73.
  35. Vadasz D, Ries V, Oertel WH. Intravenous iron sucrose for restless legs syndrome in pregnant women with low serum ferritin. *Sleep medicine* 2013; 14: 1214-6.
  36. Ghesquière L, Deruelle P, Ramdane Y, Garabedian C, Charley-Monaca C, Dalmas AF. Obstructive sleep apnea in obese pregnant women: A prospective study. *PloS one* 2020; 15: e0238733.
  37. Myoga M, Tsuji M, Tanaka R, Shibata E, Askew DJ, Aiko Y, et al. Impact of sleep duration during pregnancy on the risk of gestational diabetes in the Japan environmental and Children's study (JECS). *BMC Pregnancy Childbirth* 2019; 19: 483.
  38. Facco FL, Grobman WA, Reid KJ, Parker CB, Hunter SM, Silver RM, et al. Objectively measured short sleep

- duration and later sleep midpoint in pregnancy are associated with a higher risk of gestational diabetes. *Am J Obstet Gynecol* 2017; 217: 447.e1-.e13.
39. Pamidi S, Pinto LM, Marc I, Benedetti A, Schwartzman K, Kimoff RJ. Maternal sleep-disordered breathing and adverse pregnancy outcomes: a systematic review and metaanalysis. *Am J Obstet Gynecol* 2014; 210: 52.e1-52.e14.
40. Reutrakul S, Van Cauter E. Interactions between sleep, circadian function, and glucose metabolism: implications for risk and severity of diabetes. *Ann N Y Acad Sci* 2014; 1311: 151-73.
41. Izci-Balsarak B, Pien GW. The relationship and potential mechanistic pathways between sleep disturbances and maternal hyperglycemia. *Current diabetes reports* 2014; 14: 459.
42. Hedman C, Pohjasvaara T, Tolonen U, Suhonen-Malm AS, Myllylä VV. Effects of pregnancy on mothers' sleep. *Sleep Med* 2002; 3: 37-42.
43. Mindell JA, Jacobson BJ. Sleep disturbances during pregnancy. *J Obstet Gynecol Neonatal Nurs* 2000; 29: 590-7.
44. Ayas NT, White DP, Al-Delaimy WK, Manson JE, Stampfer MJ, Speizer FE, et al. A prospective study of self-reported sleep duration and incident diabetes in women. *Diabetes care* 2003; 26: 380-4.
45. Bassan H, Uliel-Sibony S, Katsav S, Farber M, Tauman R. Maternal Sleep Disordered Breathing and Neonatal Outcome. *Isr Med Assoc J* 2016; 18: 45-8.
46. Owusu JT, Anderson FJ, Coleman J, Oppong S, Seffah JD, Aikins A, et al. Association of maternal sleep practices with pre-eclampsia, low birth weight, and stillbirth among Ghanaian women. *Int J Gynaecol Obstet* 2013; 121: 261-5.
47. Aune D, Saugstad OD, Henriksen T, Tonstad S. Maternal body mass index and the risk of fetal death, stillbirth, and infant death: a systematic review and meta-analysis. *JAMA* 2014; 311: 1536-46.
48. Getiye Y, Fantahun M. Factors associated with perinatal mortality among public health deliveries in Addis Ababa, Ethiopia, an unmatched case control study. *BMC Pregnancy Childbirth* 2017; 17: 245.

**Citation:** Hajipour M, Soltani M, Safari-Faramani R, Khazaei S, Etemad K, Rahmani S, et al. **Maternal Sleep and Related Pregnancy Outcomes: A Multicenter Cross-Sectional Study in 11 Provinces of Iran.** *J Fam Reprod Health* 2021; 15(1): 53-60.